

Exact Groundstates of Rotating Bose Gases close to a Feshbach Resonance

N. R. Cooper

Cavendish Laboratory, Madingley Road, Cambridge, CB3 0HE, United Kingdom.

(Dated: August 14, 2003)

We study the groundstates of rotating Bose gases when interactions are affected by a nearby Feshbach resonance. We show that exact groundstates at high angular momentum can be found analytically for a general and realistic model for the resonant interactions. We identify parameter regimes where the exact groundstates are exotic fractional quantum Hall states, the excitations of which obey non-abelian exchange statistics.

PACS numbers: 03.75.Kk, 05.30.Jp, 73.43.Cd, 73.43.Nq

Dilute atomic gases are fascinating systems with which to investigate the unusual properties of Bose-condensed systems when subjected to rotation. They offer the possibility of entering a regime of high rotation rate where novel uncondensed groundstates can appear[1]. At sufficiently high angular momentum, the groundstate of a system of bosonic atoms interacting by contact repulsion is the bosonic Laughlin state[1, 2]. In addition, numerical studies [3, 4, 5] have indicated that as the angular momentum is increased many other strongly-correlated groundstates appear before the Laughlin state is reached.

In order to facilitate the experimental observation of the strongly-correlated groundstates, it has been proposed to use a Feshbach resonance to increase the energy scale of two-body interactions[6]. In addition, the Feshbach resonance induces an energy dependence to the two-body scattering amplitude and leads to a non-zero concentration of bound molecular states of the atoms[8, 9]. It is therefore possible that new strongly-correlated groundstates may form[6, 7].

In this paper we show that *exact* groundstates at high angular momentum can be found for a general and realistic model of atoms interacting via a Feshbach resonance. We show that the Laughlin state of atoms is the exact groundstate of this model over a range of parameter space. Furthermore, we identify parameter regimes where the exact groundstates are strongly-correlated atom/molecule mixtures, which are directly related to the unconventional fractional quantum Hall states proposed by Moore and Read[10] and by Read and Rezayi[11] (sometimes referred to as the “Pfaffian” and the “parafermion” states). These many-body states have very exotic physical properties, including non-abelian exchange statistics of their quasiparticle excitations[10, 11]. Exact results are extremely rare in the field of strongly-correlated many-body systems, and provide a crucial underpinning to its study. Our results establish the importance of these exotic strongly-correlated states for rotating atom gases in a realistic physical situation.

We consider a system of bosonic atoms and molecules confined to a harmonic trap with cylindrical symmetry about the z -axis. We denote the natural frequencies of the trap by $\omega_{\parallel}$ and $\omega_{\perp}$ in the axial and transverse directions. The Hamiltonian we consider is the same as that used to explore the effects of a Feshbach resonance

in other contexts[8, 9]. We write the Hamiltonian in the non-rotating frame in the form

$$\hat{H} = \hat{H}_K + \hat{H}_F + \hat{H}_I + (\hbar\omega_{\perp} + \hbar\omega_{\parallel}/2)\hat{N} + \hbar\omega_{\perp}\hat{L} \quad (1)$$

defining

$$\begin{aligned} \hat{H}_K &\equiv \int \hat{a}_{\mathbf{r}}^{\dagger} \left[\hat{h}_a - (\hbar\omega_{\perp} + \hbar\omega_{\parallel}/2) \right] \hat{a}_{\mathbf{r}} d^3\mathbf{r} \\ &\quad + \hat{m}_{\mathbf{r}}^{\dagger} \left[\hat{h}_m - (\hbar\omega_{\perp} + \hbar\omega_{\parallel}/2) \right] \hat{m}_{\mathbf{r}} d^3\mathbf{r} \\ \hat{H}_F &\equiv \Delta \int \hat{m}_{\mathbf{r}}^{\dagger} \hat{m}_{\mathbf{r}} d^3\mathbf{r} + \frac{U_{aa}}{2} \int \hat{a}_{\mathbf{r}}^{\dagger} \hat{a}_{\mathbf{r}}^{\dagger} \hat{a}_{\mathbf{r}} \hat{a}_{\mathbf{r}} d^3\mathbf{r} \\ &\quad + \frac{g}{\sqrt{2}} \int [\hat{m}_{\mathbf{r}}^{\dagger} \hat{a}_{\mathbf{r}} \hat{a}_{\mathbf{r}} + \hat{m}_{\mathbf{r}} \hat{a}_{\mathbf{r}}^{\dagger} \hat{a}_{\mathbf{r}}^{\dagger}] d^3\mathbf{r} \\ \hat{H}_I &\equiv U_{am} \int \hat{m}_{\mathbf{r}}^{\dagger} \hat{a}_{\mathbf{r}}^{\dagger} \hat{a}_{\mathbf{r}} \hat{m}_{\mathbf{r}} d^3\mathbf{r} + \frac{U_{mm}}{2} \int \hat{m}_{\mathbf{r}}^{\dagger} \hat{m}_{\mathbf{r}}^{\dagger} \hat{m}_{\mathbf{r}} \hat{m}_{\mathbf{r}} d^3\mathbf{r} \end{aligned}$$

where $\hat{a}_{\mathbf{r}}^{(\dagger)}$ and $\hat{m}_{\mathbf{r}}^{(\dagger)}$ are the (bosonic) field operators for atoms and molecules.

$\hat{H}_K$ is the kinetic energy (relative to the zero-point energy) in a frame rotating at $\omega_{\perp}$ about the $\hat{z}$ -axis, with

$$\begin{aligned} \hat{h}_a &= \frac{1}{2M} [-i\hbar\nabla - \mathbf{A}_a(\mathbf{r})]^2 + \frac{1}{2}M\omega_{\parallel}^2 z^2 \\ \hat{h}_m &= \frac{1}{4M} [-i\hbar\nabla - \mathbf{A}_m(\mathbf{r})]^2 + \frac{1}{2}(2M)\omega_{\parallel}^2 z^2 \end{aligned}$$

where $\mathbf{A}_a = M\omega_{\perp}\hat{z} \times \mathbf{r}$ and $\mathbf{A}_m = 2M\omega_{\perp}\hat{z} \times \mathbf{r}$. The eigenstates of $\hat{h}_a$ and $\hat{h}_m$ are the products of symmetric-gauge Landau level states (in the xy -coordinates) and simple harmonic oscillator states (in z). The single-particle states for atoms and molecules differ only in the natural quantum lengthscales: in the Landau level states the “magnetic lengths” for the atoms and molecules are $\ell_a = \sqrt{\hbar/(2M\omega_{\perp})}$ and $\ell_m = \ell_a/\sqrt{2}$; in the simple harmonic oscillator states, the “oscillator lengths” are $w_a = \sqrt{\hbar/(M\omega_{\parallel})}$ and $w_m = w_a/\sqrt{2}$. (In each case, the $1/\sqrt{2}$ reduction is due to the molecule/atom mass ratio of 2.)

$\hat{H}_F$ describes atom-atom scattering via the Feshbach resonance[8]: g is the strength of hybridisation; the detuning Δ is the energy difference between the lowest energy state of one molecule and the lowest energy state of two atoms in the rotating frame[12], which can be

tuned continuously with an applied magnetic field. $\hat{H}_I$ describes elastic atom-molecule and molecule-molecule scattering. We shall consider only the case of repulsive bare interactions $U_{aa}, U_{am}, U_{mm} \geq 0$ and positive detuning $\Delta > 0$. However, note that we make no assumptions concerning the relative sizes of the interactions and the level spacings of the confinement potential.

The Hamiltonian commutes with both the total number of nuclei (in atoms and molecules)

$$\hat{N} \equiv \int [\hat{a}_{\mathbf{r}}^\dagger \hat{a}_{\mathbf{r}} + 2\hat{m}_{\mathbf{r}}^\dagger \hat{m}_{\mathbf{r}}] d^3\mathbf{r}$$

and the total angular momentum (in units of $\hbar$)

$$\hat{L} \equiv \hat{\mathbf{z}} \cdot \int [\hat{a}_{\mathbf{r}}^\dagger (i\nabla \times \mathbf{r}) \hat{a}_{\mathbf{r}} + \hat{m}_{\mathbf{r}}^\dagger (i\nabla \times \mathbf{r}) \hat{m}_{\mathbf{r}}] d^3\mathbf{r}.$$

We shall look for the groundstates of $\hat{H}$ at fixed N and L . The terms in (1) proportional to N and L are then just constant energy offsets, and we need only consider the action of $\hat{H}_K$, $\hat{H}_F$ and $\hat{H}_I$.

The crucial parameter controlling the nature of the groundstates of a system of atoms at large angular momentum ($L \gg N \gg 1$) is the filling fraction[4]. In the present situation of an atom/molecule mixture, we define the *total filling fraction* in terms of N and L by

$$\nu_T \equiv \frac{N^2}{2L}.$$

Denoting the mean 2D density of nuclei in both atoms and molecules by $n_T = \bar{n}_a + 2\bar{n}_m$, where $\bar{n}_{a,m}$ are the expectation values of the atomic and molecular 2D densities, then the total filling fraction may be written

$$\nu_T = n_T(2\pi\ell_a^2) = \bar{\nu}_a + 4\bar{\nu}_m$$

where the individual filling fractions[4] for the atoms and molecules alone are $\bar{\nu}_a \equiv \bar{n}_a(2\pi\ell_a^2)$ and $\bar{\nu}_m \equiv \bar{n}_m(2\pi\ell_m^2)$. The factor of 4 appearing in this formula is a common feature of quantum Hall states containing bound pairs [13].

We now turn to look for the exact groundstates of the Hamiltonian at fixed total N and L . Substantial progress can be made by noting that, owing to the contact nature of the atom-atom interaction and the atom-molecule hybridisation, the atomic field operators in $\hat{H}_F$ can be eliminated in favour of an atom pair operator

$$\hat{P}_{\mathbf{r}}^{(\dagger)} \equiv \hat{a}_{\mathbf{r}}^{(\dagger)} \hat{a}_{\mathbf{r}}^{(\dagger)}.$$

The resulting quadratic form can be diagonalised to give

$$\begin{aligned} \hat{H}_F = & \int \left\{ \frac{\lambda_-}{\alpha^2 + \beta^2} \left(\alpha \hat{m}_{\mathbf{r}}^\dagger + \beta \hat{P}_{\mathbf{r}}^\dagger \right) \left(\alpha \hat{m}_{\mathbf{r}} + \beta \hat{P}_{\mathbf{r}} \right) \right. \\ & \left. + \frac{\lambda_+}{\alpha^2 + \beta^2} \left(\beta \hat{m}_{\mathbf{r}}^\dagger - \alpha \hat{P}_{\mathbf{r}}^\dagger \right) \left(\beta \hat{m}_{\mathbf{r}} - \alpha \hat{P}_{\mathbf{r}} \right) \right\} d^3\mathbf{r} \end{aligned}$$

where

$$\begin{aligned} \lambda_{\pm} &= \frac{1}{2} \left(\Delta + U_{aa}/2 \pm \sqrt{(\Delta - U_{aa}/2)^2 + 2g^2} \right) \\ \alpha &= g/\sqrt{2} \\ \beta &= \lambda_- - \Delta. \end{aligned}$$

$\hat{H}_F$ is positive semi-definite provided $\lambda_- \geq 0$, which is equivalent to the condition

$$\frac{g^2}{\Delta} \leq U_{aa}. \quad (2)$$

Under this condition the reduction of energy of a pair of atoms arising from hybridisation with the molecular level is less than the energy cost from the repulsive contact interaction to bring the two atoms together: there remains a net repulsion between pairs of atoms.

We shall identify exact groundstates of the Hamiltonian, under the condition (2). Our strategy is to construct many-particle states (for a given N and L) which are simultaneous zero-energy eigenstates of $\hat{H}_K$, $\hat{H}_F$, and $\hat{H}_I$. Since each of these terms is positive semi-definite, the resulting states are guaranteed to be groundstates of the total Hamiltonian (1) for given N and L .

We start from a *purely atomic state* $|\Psi\rangle_a$ (for which $\hat{m}_{\mathbf{r}}|\Psi\rangle_a = 0$), and which is an eigenstate of $\hat{N}$ and $\hat{L}$. Consider the (unnormalised) state $|\Psi\rangle$ formed by

$$|\Psi\rangle = \hat{R}|\Psi\rangle_a \quad (3)$$

$$\hat{R} \equiv \exp \left(\frac{\alpha}{\beta} \int \hat{P}_{\mathbf{r}} \hat{m}_{\mathbf{r}}^\dagger d^3\mathbf{r} \right). \quad (4)$$

Since $\hat{R}$ commutes with both $\hat{N}$ and $\hat{L}$, the number of nuclei and the total angular momentum of $|\Psi\rangle$ are both equal to those of the underlying atomic state $|\Psi\rangle_a$.

The operator $\hat{R}$ induces hybridisation of pairs of atoms in $|\Psi\rangle_a$ into molecules, in such a way that the new state (3) is annihilated by the term containing λ_+ in $\hat{H}_F$. This follows from noting that $[\hat{m}_{\mathbf{r}}, \hat{R}] = (\alpha/\beta)\hat{R}\hat{P}_{\mathbf{r}}$, so

$$\left(\beta \hat{m}_{\mathbf{r}} - \alpha \hat{P}_{\mathbf{r}} \right) \hat{R}|\Psi\rangle_a = \hat{R}(\beta \hat{m}_{\mathbf{r}})|\Psi\rangle_a = 0$$

since $\hat{m}_{\mathbf{r}}|\Psi\rangle_a = 0$. We can describe the residual effects of $\hat{H}_F$ on the states (3) by deriving the action of the operator $\hat{R}^\dagger \hat{H}_F \hat{R}$ within the subspace of purely atomic states. Using $\hat{m}_{\mathbf{r}}|\Psi\rangle_a = 0$, we find

$$\hat{R}^\dagger \hat{H}_F \hat{R} = \lambda_- \left(1 + \frac{\alpha^2}{\beta^2} \right) \int \hat{a}_{\mathbf{r}}^\dagger \hat{a}_{\mathbf{r}}^\dagger \hat{O} \hat{a}_{\mathbf{r}} \hat{a}_{\mathbf{r}} d^3\mathbf{r}. \quad (5)$$

where $\hat{O} \equiv : \exp \left(\frac{\alpha^2}{\beta^2} \int \hat{P}_{\mathbf{r}}^\dagger \hat{P}_{\mathbf{r}} d^3\mathbf{r} \right) :$ (the colons denote normal ordering). We shall refer to the operator (5) as an “effective atomic Hamiltonian”. We emphasise that it is not a unitary transformation of $\hat{H}_F$. However, its spectrum does allow one to deduce all eigenstates of $\hat{H}_F$ which are of the form (3), which is sufficient for our purposes. The residual interaction is an effective two-body




FIG. 1: Processes that generate effective (a) three-body and (b) four-body interactions between the atoms. A single (double) line denotes the propagator for an atom (molecule).

contact repulsion between the underlying atoms (with a strength reduced from $U_{aa}/2$ by the hybridisation).

Similarly, the “effective atomic Hamiltonian” from $\hat{H}_I$ is

$$\hat{R}^\dagger \hat{H}_I \hat{R} = \left(\frac{\alpha}{\beta}\right)^2 U_{am} \int [\hat{a}_\mathbf{r}^\dagger]^3 \hat{O} [\hat{a}_\mathbf{r}]^3 d^3\mathbf{r} \quad (6)$$

$$+ \left(\frac{\alpha}{\beta}\right)^4 \frac{U_{mm}}{2} \int [\hat{a}_\mathbf{r}^\dagger]^4 \hat{O} [\hat{a}_\mathbf{r}]^4 d^3\mathbf{r}. \quad (7)$$

The atom-molecule and molecule-molecule interactions generate effective three- and four-body contact interactions for the underlying atoms. These arise from the processes indicated schematically in Fig. 1.

We now consider the action of the kinetic energy operator $\hat{H}_K$ on the states (3). It is helpful to expand the field operators in the eigenstates of $\hat{h}_{a,m}$: we label these states by the quantum numbers n, m and q , where $n = 0, 1, 2, \dots$ and $m = -n, -n+1, \dots$ are the Landau level index and the angular momentum quantum number (describing motion in the xy -plane), and $q = 0, 1, 2, \dots$ is the oscillator index of the axial motion. We then define $\hat{a}_{nmq}^\dagger \equiv \int d^3\mathbf{r} \hat{a}_\mathbf{r}^\dagger \langle \mathbf{r} | n, m, q \rangle_a$, and $\hat{m}_{nmq}^\dagger \equiv \int d^3\mathbf{r} \hat{m}_\mathbf{r}^\dagger \langle \mathbf{r} | n, m, q \rangle_m$. The states $|n, m, q\rangle_m$ and the states $|n, m, q\rangle_a$ are each assumed to provide a complete basis in which arbitrary wavefunctions can be expanded (over all lengthscales for which the contact interaction Hamiltonian is assumed valid). We may then write

$$\int \hat{P}_\mathbf{r} m_\mathbf{r}^\dagger d^3\mathbf{r} = \sum_{n,m,q} \hat{P}_{nmq} m_{nmq}^\dagger \quad (8)$$

where

$$\hat{P}_{nmq}^\dagger \equiv \sum_{n_1, m_1, q_1} \sum_{n_2, m_2, q_2} \left\{ \hat{a}_{n_1 m_1 q_1}^\dagger \hat{a}_{n_2 m_2 q_2}^\dagger \int_a \langle n_1, m_1, q_1 | \mathbf{r} \rangle_a \langle n_2, m_2, q_2 | \mathbf{r} \rangle \langle \mathbf{r} | n, m, q \rangle_m d^3\mathbf{r} \right\} \quad (9)$$

We now insist that *all atoms in the state $|\Psi\rangle_a$ are in the lowest Landau level and lowest axial oscillator state*: that is, in states with $n = q = 0$. In this case, the “effective atomic Hamiltonian” $\hat{R}^\dagger \hat{H}_K \hat{R}$ vanishes identically. This surprising result follows from the fact that the $n = q = 0$ wavefunctions for a molecule, $\langle \mathbf{r} | 0, m, 0 \rangle_m$, which are proportional to

$$(x + iy)^m e^{-(x^2+y^2)/4\ell_m^2} e^{-z^2/2w_m^2}$$

provide a complete set of states for the product of any two atomic wavefunctions with $n = q = 0$, $\langle \mathbf{r} | 0, m_1, 0 \rangle_a \langle \mathbf{r} | 0, m_2, 0 \rangle_a$, which is proportional to

$$(x + iy)^{m_1+m_2} e^{-2(x^2+y^2)/4\ell_a^2} e^{-2z^2/2w_a^2}.$$

The completeness relation holds since m can span all possible values of $m_1 + m_2$ and since the arguments in the exponentials are identical due to $\ell_m = \ell_a/\sqrt{2}$ and $w_m = w_a/\sqrt{2}$. Consequently, from (8,9), the action of $\int \hat{P}_\mathbf{r} \hat{m}_\mathbf{r}^\dagger d^3\mathbf{r}$ on any state in which all atoms have $n = q = 0$ is equivalent to the action of $\sum_m \hat{P}_{0m0} \hat{m}_{0m0}^\dagger$.

In the new state (3) molecules are only in the lowest Landau level and lowest axial oscillator state ($n = q = 0$): the state (3) is therefore annihilated by $\hat{H}_K$. We emphasise that choosing to construct $|\Psi\rangle_a$ from states in which atoms are only in states with $n = q = 0$ does *not* impose further restrictions on the parameters in the model (1); we make this choice because the exact groundstates are of this form in the situations we study.

We are now in a position to deduce exact groundstates of the Hamiltonian (1) at particular values of N and L . States of the form (3) based on an underlying atomic state $|\Psi\rangle_a$ in which all atoms have $n = q = 0$ are zero-energy eigenstates of $\hat{H}_K$. To identify zero-energy eigenstates of $\hat{H}_F$ and $\hat{H}_I$, we further require that the underlying atomic wavefunction $|\Psi\rangle_a$ be annihilated by the effective two-, three-, and four-body contact interactions (5,6,7).

Zero-energy eigenstates of the two-body contact interaction (and therefore also of the three- and four-body interactions), can be constructed in the lowest Landau level for any $L \geq N(N-1)$ by choosing $|\Psi\rangle_a$ to be the bosonic Laughlin state (at $L = N(N-1)$), or the Laugh-

lin state with additional quasiholes (at $L > N(N-1)$). The Laughlin ket state of N atoms, $|\Psi_L\rangle_a$, may be constructed from its first quantised wavefunction by

$$|\Psi_L\rangle_a = \prod_{i=1}^N \int d^3\mathbf{r}_i \Psi_L(\{\mathbf{r}_i\}) \prod_{j=1}^N \hat{a}_{\mathbf{r}_j}^\dagger |0\rangle \quad (10)$$

$$\Psi_L(\{\mathbf{r}_i\}) = \prod_{i<j=1}^N (\eta_i - \eta_j)^2 e^{-\sum_i \left[\frac{|\eta_i|^2}{4\ell_a^2} + \frac{z_i^2}{2w_a^2} \right]}$$

where $\eta_i \equiv (x_i + iy_i)$ and $|0\rangle$ is the particle vacuum. (We omit normalisation factors.)

For $L = N(N-1)$ the Laughlin state is the only state in the lowest Landau level that is a zero-energy eigenstate of the two-body contact interaction. The state $|\Psi_L\rangle = \hat{R}|\Psi_L\rangle_a$ is therefore the non-degenerate groundstate of the model Hamiltonian (1) at $L = N(N-1)$, *i.e.* at total filling fraction $\nu_T = 1/2$. Note, however, that since $|\Psi_L\rangle_a$ has no amplitude for two atoms to coincide, consequently $\hat{P}_r|\Psi_L\rangle_a = 0$ and $|\Psi_L\rangle = \hat{R}|\Psi_L\rangle_a = |\Psi_L\rangle_a$. The exact groundstate at $\nu_T = 1/2$ is the bosonic Laughlin state for atoms (10). We emphasise that this result is valid for arbitrarily strong atom-atom, atom-molecule and molecule-molecule interactions. The only restrictions we impose are that $g^2/\Delta < U_{aa}$, and $\Delta, U_{aa}, U_{am}, U_{mm} \geq 0$. The pairing correlations between atoms suggested by Bhongale *et al.* [6] do not appear in the groundstate under these conditions.

It is particularly interesting to consider tuning the parameters (for example by varying the detuning Δ) such that $g^2/\Delta = U_{aa}$. Then $\lambda_- = 0$ and the effective two-body repulsion (5) vanishes: the effective atomic Hamiltonian consists only of the repulsive three- and four-body contact interactions (6,7).

Exact zero-energy eigenstates of these interactions can be constructed in the lowest Landau level for $L \geq N(N/2-1)$, from the Moore-Read (“Pfaffian”) state [10]. The Moore-Read state is the unique zero-energy eigen-

state of a three-body contact interaction at $L = N(N/2-1)$ [14]. Hence, we establish that $|\Psi_{MR}\rangle = \hat{R}|\Psi_{MR}\rangle_a$ is the non-degenerate groundstate of the Hamiltonian (1) at $L = N(N/2-1)$, which corresponds to $\nu_T = 1$ [15].

If $U_{am}/U_{mm} \ll g^2/(\Delta^2 w_a \ell_a^2)$, then the effective three-body contact interaction (6) is negligible compared to the effective four-body interaction (7) when $\nu_T \sim O(1)$. It is then of interest to set $U_{am} = 0$ and find exact groundstates of the four-body contact interaction alone (7). This can be achieved in the lowest Landau level for $L \geq N(N/3-1)$ by use of the $k=3$ Read-Rezayi (“parafermion”) wavefunction [11]. From the uniqueness of this state at $L = N(N/3-1)$ [11], we establish that $|\Psi_{k=3}\rangle = \hat{R}|\Psi_{k=3}\rangle_a$ is the non-degenerate groundstate of (1) at $L = N(N/3-1)$, corresponding to $\nu_T = 3/2$.

Note that the exact groundstates which are formed from an underlying Moore-Read state (at $\nu_T = 1$) or $k=3$ Read-Rezayi state (at $\nu_T = 3/2$) are *strongly-correlated atom/molecule mixtures* (since $\hat{P}_r|\Psi_{MR}\rangle_a \neq 0$ and $\hat{P}_r|\Psi_{k=3}\rangle_a \neq 0$). These atom/molecule mixtures will exhibit all the exotic properties that have been predicted for the Moore-Read and Read-Rezayi states of atoms alone, including non-abelian statistics for their quasiparticle excitations [10, 11]. Although these states provide exact descriptions of the groundstates only at $g^2/\Delta = U_{aa}$ (and $U_{am} = 0$ for the Read-Rezayi state), since these states have excitation gaps they should continue to describe the groundstates at $\nu_T = 1$ and $\nu_T = 3/2$ over a range of parameter space. Indeed, these states have been found to provide accurate descriptions of the groundstates of a purely atomic system ($\Delta \rightarrow \infty$) at $\nu_T = 1$ [3, 4, 5] and $\nu_T = 3/2$ [4]; it is therefore possible that they provide accurate descriptions of the groundstates at these filling fractions over the range $g^2/\Delta = U_{aa}$ to $g^2/\Delta = 0$.

I am grateful to N. Read, S. G. Bhongale and J. N. Milstein for helpful correspondence. I acknowledge support from EPSRC grant no. GR/R99027/01.

-
- [1] N. K. Wilkin, J. M. F. Gunn, and R. A. Smith, Phys. Rev. Lett. **80**, 2265 (1998).
 - [2] R. B. Laughlin, Phys. Rev. Lett. **50**, 1395 (1983).
 - [3] N. K. Wilkin and J. M. F. Gunn, Phys. Rev. Lett. **84**, 6 (2000); N. R. Cooper and N. K. Wilkin, Phys. Rev. B **60**, R16279 (1999).
 - [4] N. R. Cooper, N. K. Wilkin, and J. M. F. Gunn, Phys. Rev. Lett. **87**, 120405 (2001).
 - [5] N. Regnault and Th. Jolicoeur, Phys. Rev. Lett. **91**, 030402 (2003).
 - [6] S. G. Bhongale, J. N. Milstein, and M. J. Holland, cond-mat/0305399 (unpublished).
 - [7] U. R. Fischer, P. O. Fedichev, and A. Recati, cond-mat/0212419 (unpublished).
 - [8] E. Timmermans *et al.*, Phys. Rev. Lett. **83**, 2691 (1999).
 - [9] M. Holland, J. Park, and R. Walser, Phys. Rev. Lett. **86**, 1915 (2001).
 - [10] G. Moore and N. Read, Nucl. Phys. B **360**, 362 (1991).
 - [11] N. Read and E. H. Rezayi, Phys. Rev. B **59**, 8084 (1999).
 - [12] In terms of the detuning ϵ defined in [8] $\Delta \equiv (\epsilon - \hbar\omega_\perp - \hbar\omega_\parallel/2)$.
 - [13] B. I. Halperin, Helv. Phys. Acta **56**, 75 (1983); F. D. M. Haldane and E. H. Rezayi, Phys. Rev. Lett. **60**, 956 (1988).
 - [14] M. Greiter, X. G. Wen, and F. Wilczek, Phys. Rev. Lett. **66**, 3205 (1991).
 - [15] The Moore-Read state and $k=3$ Read-Rezayi state for atoms, $|\Psi_{MR}\rangle_a$ and $|\Psi_{k=3}\rangle_a$, can be constructed in the same way as (10) from the first quantised wavefunctions given in [10, 11].